

High time resolution observations of the January 2000 glitch in the Vela pulsar

R.G. Dodson¹ and P.M. McCulloch¹ and D.R. Lewis¹

School of Mathematics and Physics, University of Tasmania

Richard.Dodson@utas.edu.au

ABSTRACT

Pulsars are rotating neutron stars, sweeping the emission regions from the magnetic poles across our line of sight. Isolated neutron stars lose angular momentum through dipole radiation and (possibly) particle winds, hence they slow down extremely steadily, making them amongst the most reliable timing sources available. However, it is well known that younger pulsars can suffer glitches, when they suddenly deviate from their stable rotation period. On 2000 January 16 (MJD 51559) the rate of pulsation from the Vela pulsar (B0833-45) showed such a fractional period change of 3.1×10^{-6} , the largest recorded for this pulsar. The glitch was detected and reported by the Hobart radio telescope. The speedy announcement allowed the X-ray telescope, Chandra, and others, to make Target of Opportunity observations. The data placed an upper limit of 40 seconds for the transition time from the original to the new period. Four relaxation timescales are found, which are believed to be due to the variable coupling between the crust and the interior fluid. One is very short, about 60 seconds; the others have been previously reported and are 0.56, 3.33 and 19.1 days in length.

Subject headings: stars: neutron – dense matter – pulsars: individual (PSR B0833-45)

1. Introduction

Observations of pulsar glitches, in addition to providing insights into the phenomenon itself, offer one of the few probes of neutron star structure, and thus the physics of ultra-dense matter. Vela is the brightest known radio pulsar, and as it is at a declination of -45° ,

¹School of Mathematics and Physics, University of Tasmania, GPO Box 252-21, Hobart, Tasmania 7001, Australia

it is above the horizon at the Hobart Radio Observatory (Mount Pleasant) for more than 18 hours a day. It undergoes large glitches in pulse rate every few years and so provides an excellent probe of the physics of glitches.

Since 1981 the University of Tasmania has devoted a 14m diameter antenna at its Mt Pleasant Observatory to measurements of arrival times of pulses from the Vela pulsar. During this time we have observed 7 large glitches, or sudden decreases, in the period of the pulsar (McCulloch et al. 1983, 1987, 1990; McCulloch 1996).

The telescope has a single pulse observing system whose speed and sensitivity have been enhanced in order to answer a number of questions; how quickly does the crust accelerate to the new period during a glitch, how soon does the recovery from the glitch start, and what is the form of this recovery?

On 2000 January 16 (MJD 51559) the rate of pulsation jumped with a fractional period change of 3.1×10^{-6} , the largest recorded for this pulsar. The glitch was automatically detected and we issued an IAU telegram (Dodson et al. 2000) within 12 hours, allowing the X-ray telescope, Chandra, to make Target of Opportunity (TOO) observations (Helfand et al. 2001). These observations have so far failed to find the signature of neutron star heating, which was the driver for the TOO, but have produced spectacular images of the X-ray pulsar wind nebula.

2. Observations

The reported observations were made simultaneously at 635 MHz, 990 MHz and 1390 MHz, to allow continuous measurement of the dispersion measure (DM). Dual channel receivers with an equivalent system temperature of 60 K are used at each frequency, except for 1390 MHz, where only the right handed circular polarisation is observed. Receiver bandwidths of 250 kHz at 635 MHz, 800 kHz at 990 MHz and 2 MHz at 1390 MHz limit the pulse broadening from interstellar dispersion to less than 1% of the pulse period. The detected output from each receiver was folded for 2 minutes to give integrated pulse profiles from 1,344 pulses. The profiles were subsequently combined to give profiles of total intensity. The signal-to-noise ratio in each total intensity profile is typically 30:1, allowing a mean pulse arrival time to be determined to an accuracy of $80\mu\text{s}$ at 635 MHz, $60\mu\text{s}$ at 990 MHz and $180\mu\text{s}$ at 1390 MHz per integration.

The recent improvements in the time resolution have been achieved by incoherently de-dispersing over 8 adjacent channels at 990 MHz, thereby increasing the signal to noise ratio and allowing observations of single pulses. The de-dispersed bandpass is sampled at 2 kHz

and recorded directly onto disk for later retrieval, while an ‘on the fly’ monitoring system folds on a ten second basis for which the RMS is $85\mu\text{s}$. These arrival times are monitored and if a glitch is detected a warning is issued and the single pulse data are retained.

3. Timing fits

The full, folded, timing profiles are stored in EPN format (Lorimer et al. 1998), cross correlated with high signal to noise profile templates, then fitted to find the time of arrival (TOA) of the pulse.

A Taylor expansion is parameterised in terms of the spin-down model phase (ϕ) around a selected epoch (t relative to t_0) with frequency ν , and frequency derivative $\dot{\nu}$, and has the form;

$$\phi_m(t) = \phi_0 + \nu(t - t_0) + \frac{1}{2}\dot{\nu}(t - t_0)^2 + \frac{1}{6}\ddot{\nu}(t - t_0)^3 \quad (1)$$

The post glitch frequency change can be written as the sum of the pre-glitch frequency plus the permanent change at the time of the glitch, plus other decay terms:

$$\nu(t) = \nu_0 + \Delta\nu_p + \Delta\dot{\nu}_p t + \frac{1}{2}\Delta\ddot{\nu}_p t^2 + \sum_n \Delta\nu_n e^{-t/\tau_n} \quad (2)$$

However, it should be noted that some investigators set the permanent change terms, $\Delta\dot{\nu}_p$ and $\Delta\ddot{\nu}_p$ to zero. We have allowed $\Delta\dot{\nu}_p$ to take non-zero values, but keep $\Delta\ddot{\nu}_p$ as zero. Both of these terms affect only the longest time-scales, where both timing noise and the occurrence of the subsequent glitches make definitive separation impossible.

The data presented here were recorded between MJD 51505 and 51650. The arrival time data have been transformed to the Solar System barycentre using standard techniques. The position and proper motion of the Vela pulsar was defined by data from the Radio VLBI position of Legge (2001). The recorded TOAs from all frequencies and both systems were fitted in the program **TEMPO**². The results of this fit are given in Table 1.

Figure 1a shows the residuals from the pre-jump fit for data taken on 2000 January 16 (MJD 51559) with the full polarisation system. Shortly after 07:34 UT, the residuals diverge

²See <http://pulsar.princeton.edu/tempo> (Taylor, Manchester, Nice, Weisberg, Irwin, Wex, & Standish 1970)

from the fit, indicating a sudden decrease in pulse period. Figure 1b shows an hour of data starting at MJD 51559.3. Individual data points represent ten second averages constructed from the single pulse data. The period jump occurs on a very short timescale, without warning. The observations are consistent with an instantaneous change in period; modeling has shown that a spin-up timescale of forty seconds would produce a three sigma signal.

The separation into four time-scales is clear. The longer three decay terms are similar to those previously reported (Alpar et al. 1993; Flanagan 1990), and are in an approximately equal ratio of 5.9:5.7. These have been associated with the vortex creep models by Alpar et al. (1993) and others. The fast decay timescale, not previously observed (or observable) is shown separated from the other effects in figure 2. We have subtracted the terms found by **TEMPO** in the 2 minute data from the single pulse data folded for 10 seconds. In this plot a gradual spin-up (as opposed to instantaneous) would be a negative excursion around the projected time of the glitch, as we’d have overestimated the phase in the model. We see a positive excursion, followed by a rapid decay. A positive excursion could be produced by the pulsar slowing down just before the glitch or, more likely, if the estimate of the glitch epoch was too early because there was an extra component not resolvable in the 2 minute data. Once the original fit was removed this would give a linear rise with the gradient $\Delta\nu$ found in the 2 minutes data, followed by a decay. We have fitted this rise ($\Delta\nu\Delta t$) followed by a forth exponential decay term to give the true glitch epoch and the fastest decay term.

4. The single pulse system

Since the acceleration of the crust cannot be instantaneous, it should be possible to observe the spin-up of the rotation period of the pulsar. The parallel single pulse system designed to observe this has a three σ detection limit of less than 40 seconds. This is in contrast to the observations made on the Crab (Wong et al. 2001), where the spin-up timescale has been observed to take about half a day. No variation could be seen either before or after the glitch in profile shape, magnitude, or polarisation in the radio observations on any time-scale. The raw time series using both long and short timescales has been closely examined to look for other periodic and quasi-periodic signals, but no extra terms were found.

5. Estimates for the force from the vortex lines

We will use our data to explore one model of pulsar glitches in which the glitches are a consequence of pinning of superfluid vortices to the stellar crust. As the neutron star slows due the external drag on the crust, an angular momentum excess develops in the pinned superfluid. Ultimately the vortices unpin catastrophically, transferring the excess momentum to the crust, producing a glitch.

Our high time resolution studies permit constraints on the manner in which the liquid interior imparts its angular momentum to the crust. This coupling is thought to be due to dissipative motion of vortices through the crustal lattice after unpinning. Various papers have tackled calculation of the drag on vortex lines within the superfluid (e.g. Bildsten & Epstein (1989); Epstein & Baym (1992); Jones (1992)) produced by vortex-neutron scattering in the crust. We will ignore the details of the interaction here, as it is not a simple process. However, given that we have a lower limit for the spin-up time, we can obtain an observational limit for the force per length, f , that a vortex line exerts on the superfluid during this spin-up phase in Vela. Assuming that pinned vorticity in the crust is responsible for the glitch, it is straightforward to calculate the torque required to spin-up the pulsar, and then, by assuming canonical neutron star radii, crust depths and vortice lengths, give the force per unit length.

During the spin-up, the torque on the crust, $\mathcal{N}$, is:

$$\mathcal{N} = I_s \Delta\Omega_s / \Delta t$$

where I_s is the moment of inertia of the portion of the superfluid that drives the glitch and $\Delta\Omega_s$ is the change in angular velocity of that component, Δt is the spin-up time. Since angular momentum is conserved in the glitch, the torque is

$$\mathcal{N} = I_{crust} \Delta\Omega / \Delta t$$

where I_{crust} is the moment of inertia of the solid crust plus any component that is coupled to it over a timescale in less than the spin-up time Δt , and $\Delta\Omega$ is the observed spin-up of the crust over Δt .

In magnitude, the torque is

$$\mathcal{N} = F_\phi R$$

where F_ϕ is the azimuthal component of the force that vortices exert on the lattice during

the glitch, and R is the average distance of vortices from the rotation axis, approximately the stellar radius. The azimuthal force is

$$F_\phi = f H N_v$$

where f is the force exerted per unit length per vortex, H is the average length of a vortex, and N_v is the total number of vortices that are participating in the spin-up.

The number of vortices is

$$N_v = n_v 2\pi R \Delta R$$

where n_v is the number of vortices per unit area and ΔR is the thickness of the pinning region (approximately the thickness of the crust). The number density of vortices in a rotating superfluid is

$$n_v = 2\Omega_s / \kappa$$

where Ω_s is the superfluid rotational velocity, approximately equal to the rotational velocity of the crust and κ is the quantum of circulation of a neutron fluid (0.0019 cm²/s). Putting these expressions together gives a constraint for f :

$$f = I_{crust} \frac{\kappa \Delta \Omega / \Delta t}{H 4\pi \Omega R^2 \Delta R} \quad (3)$$

ΔR , the thickness of the crust, is about 0.1 km. We take H , the length of a vortex, to be roughly ΔR .

We defined I_{crust} as the moment of inertia of the solid plus any other component coupled to it in a timescale less than Δt . If the core fluid is tightly coupled to the solid crust over Δt , then I_{crust} is approximately the moment of inertia of the entire star (about 10⁴⁵ g cm²). On the other hand, if the core is decoupled over Δt then I_{crust} would correspond to the moment of inertia of the solid crust only, which is about 10⁴³ g cm² for a moderate equation of state. So we can obtain a lower limit on f .

$$f \geq 10^{12} I_{crust,43} \text{ dyn/cm}$$

Where $I_{crust,43}$ is the crust inertia in units of 10⁴³ g cm². Vortices could pin to the lattice with a force $\sim 10^4$ times this (Link & Cutler 2002), but our observations constrain the value

of f during the dissipative regime, whilst the pulsar is being spun up. Putting the same parameters into the Crab time-resolved glitch of MJD 50259.93 (Wong et al. 2001) we find that in that case $f \approx 4 \times 10^5$ dyn/cm. This is further evidence, if more was needed, that the Crab pulsar is a very different object from the Vela pulsar. The major difference between the Crab and the Vela pulsars is the age, and therefore the temperature. Jones (1992) found a strong temperature dependence for some scattering processes, and these differences may be key to our understanding of the momentum transfer in glitches. Further spin-up observations for both the Crab and Vela are required.

6. Future development

Further improvements of the single pulse system are being undertaken, with a new coherent dedispersion system, which will allow a detection limit of a few seconds, the order of the fastest coupling times in all theoretical models. This should also allow us to measure f , the force per unit length on the vortices, and confirm the fast decay term. Higher time resolution will allow further constraints on the coupling of the crust to the liquid interior, including the core. We will also co-ordinate with Chandra to reduce the TOO delay time in order to determine whether a very short lived thermal pulse might be emitted.

The day to day maintenance of the 14m telescope, over 20 years has been a herculean task, to which too many people to name have contributed. Also we greatly appreciated the extremely helpful suggestions on the text from the referee, B. Link and also from S. Johnston.

REFERENCES

- Alpar, M. A., Chau, H. F., Cheng, K. S., & Pines, D. 1993, *Astrophys. J.*, 409, 345
- Bildsten, L. & Epstein, R. I. 1989, *Astrophys. J.*, 342, 951
- Dodson, R. G., McCulloch, P. M., & Costa, M. E. 2000, *IAU Circ.*, 7347, 2
- Epstein, R. I. & Baym, G. 1992, *ApJ*, 387, 276

- Flanagan, C. S. 1990, *Nature*, 345, 416
- Helfand, D. J., Gotthelf, E. V., & Halpern, J. P. 2001, *ApJ*, 556, 380
- Jones, P. B. 1992, *MNRAS*, 257, 501
- Legge, D. 2001, PhD thesis, School of Maths and Physics, University of Tasmania
- Link, B. & Cutler, C. 2002, to be published in *MNRAS*
- Lorimer, D. R., Jessner, A., Seiradakis, J. H., Lyne, A. G., D’Amico, N., Athanasopoulos, A., Xilouris, K. M., Kramer, M., & Wielebinski, R. 1998, *Astr. Astrophys. Suppl.*, 128, 541
- McCulloch, P. M. 1996, in *ASP Conf. Ser. 105: IAU Colloq. 160: Pulsars: Problems and Progress*, 97
- McCulloch, P. M., Hamilton, P. A., McConnell, D., & King, E. A. 1990, *Nature*, 346, 822
- McCulloch, P. M., Hamilton, P. A., Royle, G. W. R., & Manchester, R. N. 1983, *Nature*, 302, 319
- McCulloch, P. M., Klekociuk, A. R., Hamilton, P. A., & Royle, G. W. R. 1987, *Australian Journal of Physics*, 40, 725
- Taylor, J., Manchester, R., Nice, D., Weisberg, J., Irwin, A., Wex, N., & Standish, E. 1970
- Wong, T., Backer, D. C., & Lyne, A. G. 2001, *Astrophys. J.*, 548, 447

Parameters for Epoch 51559		
ν/Hz	$\dot{\nu}/Hz\ s^{-1}$	$\ddot{\nu}/Hz\ s^{-2}$
11.194615396005	-1.55615E-11	1.028E-21
$\Delta\nu_p/Hz$	$\Delta\dot{\nu}_p/Hz\ s^{-1}$	
3.45435(5)E-05	-1.0482(2)E-13	
τ_n	$\Delta\nu_n/10^{-6}Hz$	
1.2 $\pm$ 0.2mins	0.020(5)	
00.53(3) days	0.31(2)	
03.29(3) days	0.193(2)	
19.07(2) days	0.2362(2)	
DM	67.99	

Table 1: Parameters for the glitch epoch 51559.3190. Errors are the one sigma values. The data fit is from MJD 51505 through to 51650 (November 1999 to April 2000).




Fig. 1.— Arrival time residuals from the pre-glitch model for both the three frequency (1a) and the single pulse (1b) systems. Figure 1a has folds of two minutes and figure 1b has folds of ten seconds.




Fig. 2.— The previously unobserved fast decay, with ten second folds. The other decay terms are removed revealing the later start, and decay of the fastest term.